


UNITED STATES



ADOLESCENT AND ADULT USE (Ages 12 and Older)

Percent (%) Currently Using Drug

	2001	2000	1999	1998	1997	1996	1995	1994	1993	1992	1991	1990
Alcohol	48.3	46.6	46.4	51.7	51.4	51.0	52.2	53.9	50.8	49.0	52.2	52.6
Binge Alcohol Use *	20.5	20.6	20.2	15.6	15.3	15.5	15.8	16.5	14.6	14.5	15.5	14.4
Heavy Alcohol Use **	5.7	5.6	5.7	5.9	5.4	5.4						
Any Tobacco	29.5	29.3	30.2									
Cigarettes	24.9	24.9	25.8	27.7	29.6	28.9	28.8	28.6	29.6	31.9	33.0	32.6
Smokeless Tobacco	3.2	3.4	3.4	3.1	3.2	3.2	3.3	3.3	3.2	4.0	3.7	3.9
Any Illicit Drug	7.1	6.3	6.3	6.2	6.4	6.1	6.1	6.0	5.9	5.8	6.6	6.7
Marijuana/Hashish	5.4	4.8	4.7	5.0	5.1	4.7	4.7	4.8	4.6	4.7	5.1	5.4
Inhalant	0.2	0.3	0.3	0.3	0.4	0.4	0.4	0.4	0.3	0.3	0.4	0.4
Hallucinogen	0.6	0.4	0.4	0.7	0.8	0.6	0.7	0.5	0.4	0.4	0.5	0.4
LSD	0.1	0.2	0.2	0.3	0.2	0.2	0.3	0.2				
PCP	0.0	0.0	0.0	0.0	0.2	0.1	0.0	0.0	0.0	0.0	0.0	
Ecstasy	0.3											
Cocaine (any form)	0.7	0.5	0.7	0.8	0.7	0.8	0.7	0.7	0.7	1.0	0.9	
Crack Cocaine	0.2	0.1	0.2	0.2	0.3	0.3	0.2	0.2	0.3	0.2	0.3	0.3
Heroin	0.1	0.1	0.1	0.1	0.2	0.1	0.1	0.1	0.0	0.0	0.0	0.0
Pain Reliever (non-medical use)	1.6	1.2	1.2	0.8	0.7	0.9	0.6	0.7	0.8	0.9	0.8	0.9
Stimulant (non-medical use)	0.5	0.4	0.4	0.3	0.3	0.4	0.4	0.3	0.5	0.3	0.4	0.6
Methamphetamine	0.3	0.2	0.2									
Sedative (non-medical use)	0.1	0.1	0.1	0.1	0.1	0.1	0.2	0.1	0.2	0.2	0.2	0.2
Tranquilizer (non-medical use)	0.6	0.4	0.5	0.3	0.4	0.4	0.4	0.5	0.6	0.8	1.1	0.6

Percent (%) Ever Using Drug

Alcohol	81.7	81.0	81.3	81.3	81.9	82.6	82.3	84.2	82.6	81.9	83.6	82.2
Any Tobacco	71.4	70.5	72.0									
Cigarettes	67.2	66.5	68.2	69.7	70.5	71.6	71.8	73.3	73.3	73.1	74.9	75.4
Smokeless Tobacco	19.1	18.5	19.1	17.2	17.3	17.0	17.0	17.2	15.9	18.2	17.5	17.5
Any Illicit Drug	41.7	38.9	39.7	35.8	35.6	34.8	34.2	34.4	34.2	33.3	34.1	34.2
Marijuana/Hashish	36.9	34.2	34.6	33.0	32.9	32.0	31.0	31.1	31.0	30.2	30.5	30.5
Inhalant	8.1	7.5	7.8	5.8	5.7	5.6	5.7	5.8	5.9	5.3	6.1	5.7
Hallucinogen	12.5	11.7	11.3	9.9	9.6	9.7	9.5	8.7	9.0	8.3	8.4	7.9
LSD	9.0	8.8	8.7	7.9	7.8	7.7	7.5	7.0	7.0	6.7	6.4	5.8
PCP	2.7	2.6	2.6	3.5	3.0	3.2	3.2	2.8	2.7	2.7	2.4	2.0
Ecstasy	3.6	2.9										
Cocaine (any form)	12.3	11.2	11.5	10.6	10.5	10.3	10.3	10.4	11.3	10.9	11.5	11.2
Crack Cocaine	2.8	2.4	2.7	2.0	1.9	2.2	1.8	1.9	1.9	1.5	2.1	1.5
Heroin	1.4	1.2	1.4	1.1	0.9	1.1	1.2	1.0	1.0	0.8	1.2	0.8
Pain Reliever (non-medical use)	9.8	8.6	9	5.3	4.9	5.5	6.1	6.0	6.4	6.1	6.8	6.3
Stimulant (non-medical use)	7.1	6.6	7.2	4.4	4.5	4.7	4.9	4.6	4.8	5.0	5.6	5.5
Methamphetamine	4.3	4.0	4.3	2.1	2.5	2.3	2.2					
Sedative (non-medical use)	3.3	3.2	3.5	2.1	1.9	2.3	2.7	2.6	2.6	2.6	3.2	2.8
Tranquilizer (non-medical use)	6.2	5.8	6.3	3.5	3.2	3.6	3.9	4.0	4.2	4.7	5.1	4.0
Steroid								0.3	0.4	0.3	0.5	
Needle Drug Use	1.5							1.4	1.4	1.5	1.9	1.7

=> Beginning in 1999, data are not directly comparable to earlier years due to use of a computer-assisted instrument that replaced the paper-and-pencil surveys.

* "Binge" alcohol use in the National Household Survey is defined as consuming 5 or more drinks on the same occasion on at least one day in the past 30 days.

** "Heavy" alcohol use is defined as consuming 5 or more drinks on the same occasion on each of 5 or more days in the past 30 days.